

The
National
Breast Cancer *Audit*



Report on New Zealand episodes diagnosed in 2010

Report prepared for:
BreastScreen Aotearoa
December 2011

Prepared by:



Australian Safety and Efficacy Register of New Interventional Procedures – Surgical

of



Royal Australasian College of Surgeons

on behalf of:



Contact details

199 Ward St

North Adelaide, SA 5006

Phone (08) 8219 0900

Fax (08) 8219 0999

Email breast.audit@surgeons.org

Web <http://www.surgeons.org/nbca>

Table of contents

Acknowledgments	5
Funding	5
Introduction.....	6
1. Background Information.....	7
1a: Referral source for the New Zealand episodes	7
1b. Invasive and DCIS episodes by referral source [†]	8
1c. Private and public status of the episodes by referral source	9
1d. Age of the patients by referral source	10
1e. Gender of the patients by referral source	10
2. Invasive Tumour Characteristics	11
2a: Type of invasive tumour by referral source [†]	11
2b: Size of invasive tumour by referral source [†]	12
2c: Histological grade of invasive tumour by referral source [†]	13
2d: Lymphatic vascular invasion of invasive tumour by referral source [†]	14
2e: Bilateral synchronous status of invasive tumour by referral source [†]	14
2f: Menopausal status for invasive tumour by referral source [†]	15
2g: Oestrogen receptor status of invasive tumour by referral source [†]	16
2h: Progesterone receptor status of invasive tumour by referral source [†]	16
2i: HER2 Receptor status of invasive tumour by referral source [†]	17
3. DCIS Tumour Characteristics.....	18
3a: Size of DCIS tumours by referral source [†]	18
3b: Histological grade of DCIS tumour by referral source [†]	18
3c: Necrosis of DCIS tumour by referral status [†]	19
3d: Bilateral synchronous status of DCIS tumours by referral status [†]	19
3e: Menopausal status for the DCIS tumours by referral source [†]	20
4. Breast Surgery Treatment	21
4a: First breast surgery performed for invasive cancer by referral source [†]	21
4b: Further breast surgery after breast conserving surgery for invasive cancer by referral source [†]	21
4c: Reconstruction after mastectomy for invasive cancer by referral source [†]	21
4d: First breast surgery performed for DCIS by referral source [†]	22
4e: Further surgery after breast conserving surgery for DCIS by referral source [†]	22
4f: Reconstruction performed after mastectomy for DCIS by referral source [†]	22
5. Axillary surgery treatment	24
5a. Axillary procedures for invasive cancer by referral source [†]	24
5b. Axillary procedures for ≤3cm invasive cancer by referral source [†]	25
5c. Axillary procedures for >3cm invasive cancer by referral source by referral source.....	26
5d. Axillary procedures for DCIS treated with breast conserving surgery only by referral source [†]	27
5e. Axillary procedures for DCIS treated with mastectomy by referral source [†]	28
6. Margins of excision for breast surgery	29
6a. Margins of excision for invasive cancer by referral source [†]	29
6b. Margins of excision for DCIS cancer by referral source [†]	29

7. Radiotherapy treatment.....	30
7a: Radiotherapy for invasive cancer treated with breast conserving surgery by referral source [†]	30
7b: Radiotherapy for invasive cancer which had mastectomy by referral source	30
7c: Radiotherapy for high risk invasive cancer treated with mastectomy by referral source	31
7d: Radiotherapy for DCIS treated with breast conserving surgery by referral source [†]	31
7e: Radiotherapy for DCIS cancer which had mastectomy by referral source [†]	31
8. Hormonal treatment	32
8a. Hormonal treatment for oestrogen positive invasive tumours by referral source [†]	32
8b. Hormonal treatment for oestrogen negative invasive cancer by referral source [†]	34
9. Chemotherapy treatment.....	36
9a. Chemotherapy treatment for invasive cancer for ≤ 70 years old patients by referral source [†]	36
9b. Chemotherapy treatment for invasive cancer for > 70 years old patients by referral source [†]	37
10. Herceptin (Trastuzumab) treatment	38
10a. Herceptin (Trastuzumab) treatment for >1 cm HER2 positive OR node positive HER2 positive invasive cancer by referral source [†]	38
11. Summary.....	39
11a. Significant differences in invasive tumour characteristics and breast cancer treatments between BSA and not BSA patients	40
11b. DCIS tumour characteristics and breast cancer treatments between BSA and not BSA patients	41
References.....	42

Acknowledgments

This report was developed by the National Breast Cancer Audit. The audit is directed by the Breast Surgeons of Australia and New Zealand and managed by the Royal Australasian College of Surgeons Research, Audit and Academic Surgery Division and Australian Safety and Efficacy Register of New Interventional Procedures – Surgical (ASERNIP-S).

The data analysis and development of the report was undertaken by Dr Primali de Silva.

Funding

Funding for the data analysis and development of the report was provided by the Ministry of Health New Zealand, through BreastScreen Aotearoa.

Introduction

The National Breast Cancer Audit (NBCA) was initiated in 1998 and collects data on the surgical care of early breast cancer patients in Australia and New Zealand. The audit is managed by the Australian Safety and Efficacy Register of New Interventional Procedures – Surgical (ASERNIP-S) Program, being part of the Research, Audit and Academic Surgery Division of the Royal Australasian College of Surgeons.

A Structured Query Language (SQL) query has been written to extract New Zealand data with a diagnosis date of 2010 (if diagnosis date was not provided, first surgery date was used) from the restored NBCA online database on 07 September 2011. In addition to this, a data set from Auckland Breast Cancer Register for January to June 2010 was also included in the report. This dataset was scheduled to be uploaded into the NBCA database through the institutional upload program later in 2011.

There were 9359 cases reported to the NBCA in 2010 and 2126 of these cases were from New Zealand. Out of the 271 surgeons who contributed to the audit in 2010, 72 were from New Zealand.

In the report, percentage case volumes for New Zealand data have been reported by referral source under the following main headings:

1. Background Information
2. Invasive Tumour Characteristics
3. DCIS Tumour Characteristics
4. Breast Surgery Treatment
5. Axillary Surgery Treatment
6. Margins of excision for breast surgery
7. Radiotherapy Treatment
8. Hormonal Treatment
9. Chemotherapy Treatment
10. Herceptin (Trastuzumab) Treatment

In some of the treatment sections, the relevant guidelines and/or NBCA Key Performance Indicators (KPIs) have been listed. The reader can clearly see the percentage of cases which follow the guidelines and NBCA KPIs from this data.

The number of cases reported from Breast Screen Aotearoa (BSA) and other referral sources for each category were compared using chi-square tests via the Statistical Package for Social Sciences software (SPSS Inc., Chicago, IL, USA). A statistical significance level of $P < 0.05$ was used. (P value was not calculated if the number of observations per category was zero.)

Background information, tumour characteristics and breast cancer treatments that are significantly different between “BSA” and “not BSA” referral are listed in the summary section.

Definitions of the terms provided in the report are from the National Breast Cancer Audit Data Dictionary.

In this report, “Unknown”, “Not yet” and missing data are reported as “not known”.

1. Background Information

1a: Referral source for the New Zealand episodes

Referral source	Percentage
Breast Screen Aotearoa (BSA) (N=824)	38.76%
Not BSA (N=1295) - Symptomatic from GP (N=1095)	51.51%
Breast Screen Australia (N=6)	0.28%
Other (N=194)	9.13%
Not known (N=7)	0.33%
Total (N=2126)	100%

Comments:

While the majority of breast cancer cases from the New Zealand were referred as symptomatic from a GP (52%), BSA was the second most common referral source (39%).

Audit data used:

Information is derived from the audit question “referral source” which allows the options of symptomatic from General Practitioner (GP), Breast Screen Australia, Breast Screen Aotearoa (NZ) and Other.

Definitions:

Referral source records the source from which the person was referred to the surgeon. Symptomatic patients are referred to a breast surgeon when presenting to a GP or other physician with symptoms such as a breast lump, pain, or discharge. Patients referred from “Other” sources may include private screening programs.

1b. Invasive and DCIS episodes by referral source[†]

Referral source	Invasive breast cancer	DCIS
BSA (N=824)	79.85%	20.15%
Not BSA (N=1294)	93.20%	6.80%
P value	< 0.001	< 0.001

[†]Referral source was not known for 7 records and invasive/in situ status was not known for 1 Not BSA patient.

Comments:

The majority of the New Zealand breast cancer episodes were invasive (88%). There was a significantly higher percentage of DCIS cases in the BSA referral group (20%) than in the not BSA referral groups (7%).

Audit data used:

Information is derived from the audit question “invasive/in situ cancer”.

Definitions:

- Invasive – a cancer which has grown beyond its site of origin and invaded neighbouring tissue.
- DCIS – the presence of any malignant tumour which has not yet become invasive but is confined to the layer of cells from which it arose. A form of pre-invasive cancer.

1c. Private and public status of the episodes by referral source[†]

Referral source	Private	Public
BSA (N=824)	25.61%	74.39%
Not BSA (N=1295)	28.89%	71.12%
P value	0.100	0.100

[†]Referral source was not known for 7 records.

Comments:

The majority of the New Zealand patients were public (72%). The percentages of public and private patients were not significantly different between the BSA group and not BSA group.

Audit data used:

Information is derived from the audit question “public/private” which allows the options of private and public.

Definitions:

Public – a person, eligible for public healthcare who on admission to a recognised hospital or soon after:

- receives a public hospital service free of charge or
- elects to be a public patient or
- whose treatment is contracted to a public hospital

Private – a person who, on admission to a recognised hospital or soon after:

- elects to be a private patient treated by a medical practitioner of his or her choice
- elects to occupy a bed in a single room (where such an election is made, the patient is responsible for meeting certain hospital charges as well as the professional charges raised by treating medical practitioner) or
- a person, eligible for public healthcare, who chooses to be admitted to a private hospital

1d. Age of the patients by referral source[†]

Referral source	≤40 years	41-50 years	51-60 years	61-70 years	>70 years
BSA (N=824)	0.12%	20.75%	36.17%	40.29%	2.67%
Not BSA (N=1295)	8.88%	24.09%	21.70%	14.75%	30.58%
P value	<0.001	0.074	<0.001	<0.001	<0.001

[†]Referral source was not known for 7 records.

Comments:

The percentage of 51-70 year old patients was significantly higher in the BSA group (76%) than in the not BSA group (36%). The percentage of younger patients (< 40 years) and older patients (>70 years) was significantly lower in the BSA group (0.1%, 3%) than in the not BSA group (9%, 31%). The percentage of patients in the 41-50 years group was not significantly different between the BSA and not BSA groups.

Audit data used:

Information is derived from a calculation using audit questions “diagnosis date” and “date of birth”. (If diagnosis date was not available, the first surgery date was used.)

Definitions:

Diagnosis date: The date upon which the cancer diagnosis was made

Surgery date: The date upon which breast cancer surgery was done

Date of birth: Patient’s date of birth

1e. Gender of the patients by referral source[†]

Referral source	Female	Male
BSA (N=824)	100.00%	0.00%
Not BSA (N=1295)	99.00%	1.00%
P value	Not calculated	Not calculated

[†]Referral source was not known for 7 records.

Comments:

Only 1% of the New Zealand patients were males and none of the male patients were referred from BSA for the treatments.

Audit data used:

Information is derived from the audit question “gender” which allows the options of female and male.

Definitions:

Female: female patient; Male: male patient

2. Invasive Tumour Characteristics

2a: Type of invasive tumour by referral source[†]

Referral source	1	2	3	4	5	6	7	8
BSA (N= 649)	79.82%	10.32%	1.69%	1.85%	3.39%	0.31%	1.85%	0.77%
Not BSA (N=1160)	80.95%	9.74%	1.64%	2.24%	1.03%	0.52%	2.41%	1.47%
P value	0.559	0.692	0.927	0.577	<0.001	0.520	0.433	0.196

[†]Referral source was not known for 4 invasive records. Tumour types were not known for 9 BSA and 46 not BSA patients.

1 – Ductal Carcinoma Not Otherwise Specified (NOS); 2 - Invasive Lobular; 3 – Other invasive of mixed type; 4 – Other Neoplasm; 5 – Tubular; 6 – Medullary; 7 – Mucinous; 8- Basal like;

Comments:

Most (81%) of the New Zealand invasive tumours were Ductal Carcinoma NOS. The percentages of Tubular (3%) were significantly higher in the BSA group than in the not BSA group (1%). The percentages of other tumours were not significantly different between BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “invasive histological type of tumour” which allows the options of ductal carcinoma NOS, invasive lobular, tubular, medullary, mucinous, other invasive of mixed type, other neoplasm, basal-like.

Definitions:

Tumour type defines the microscopic appearance of the invasive breast cancer cells in the principal tumour.

2b: Size of invasive tumour by referral source[†]

Referral source	<10 mm	10-14 mm	15-19 mm	20-29 mm	30-39 mm	≥40 mm
BSA (N= 649)	28.66%	27.89%	19.88%	14.64%	4.47%	4.47%
Not BSA (N=1133)	10.15%	13.95%	16.33%	29.66%	13.50%	16.42%
P value	<0.001	<0.001	0.059	<0.001	<0.001	<0.001

[†] Referral source was not known for 4 invasive records. Invasive tumour size was not known for 9 BSA and 73 not BSA patients.

Comments:

The percentage of patients with smaller tumours (< 15 mm) was significantly higher in the BSA group (57%) than in the not BSA group (24%). The percentage of patients with larger tumours (>20 mm) was significantly higher in not BSA group (60%) than in the BSA group (24%). The percentage of patients with 15-19 mm invasive tumours was not significantly different between BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “invasive tumour size in mm”.

Definitions:

Tumour size refers to the maximum diameter in millimetres of the furthest points of extension of the invasive tumour cells in the principal tumour.

2c: Histological grade of invasive tumour by referral source[†]

Referral source	Grade 1	Grade 2	Grade 3
BSA (N= 640)	34.84%	46.09%	19.06%
Not BSA (N=1138)	17.75%	42.09%	40.16%
P value	< 0.001	0.102	< 0.001

[†] Referral source was not known for 4 invasive records. Histological grade of the invasive tumours were not known for 18 BSA and 68 not BSA patients.

Comments:

The percentage of patients with Grade 1 tumours was significantly higher in the BSA group (35%) than in the not BSA group (18%). The percentage of patients with Grade 3 tumours was significantly higher in the not BSA group (40%) than in the BSA group (19%). There was no significant difference for the Grade 2 invasive tumours between BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “invasive histological grade of tumour” which allows the options of grade 1, grade 2, and grade 3.

Definitions:

Histological grade is the degree of differentiation of the breast cancer or the degree to which it resembles normal tissue as assessed by the pathologist according to Pathology Reporting Guidelines. The histological grade is calculated by adding three scores (mitosis score, nuclear score and tubular differentiation score):

Grade 1 – Total score of 3-5

Grade 2 – Total score of 6-7

Grade 3 – Total score of 8-9

2d: Lymphatic vascular invasion of invasive tumour by referral source[†]

Referral source	Present	Absent
BSA (N= 640)	14.06%	85.94%
Not BSA (N=1153)	33.74%	66.26%
P value	< 0.001	< 0.001

[†] Referral source was not known for 4 invasive records. Lymphatic vascular invasion was not known for 18 BSA and 53 not BSA patients.

Comments:

In the majority (74%) of the New Zealand patients, lymphatic vascular invasion was absent. The percentage of patients with vascular lymphatic invasion was significantly lower in the BSA group (14%) than in the not BSA group (34%).

Audit data used:

Information is derived from the audit question “vascular/lymphatic invasion” which allows the options of present and absent.

Definitions:

Lymphatic vascular invasion present - tumour cells observed within the lumen of blood or lymphatic vessels.

2e: Bilateral synchronous status of invasive tumour by referral source[†]

Referral source	Bilateral synchronous	Not bilateral synchronous
BSA (N= 656)	2.44%	97.56%
Not BSA (N=1205)	3.40%	96.60%
P value	0.249	0.249

[†] Referral source was not known for 4 invasive records. Bilateral synchronous status for invasive tumours was not known for 2 BSA and 1 not BSA patients.

Comments:

Most (97%) of the New Zealand invasive cancers were not bilateral synchronous and there was no significant difference in the percentage of bilateral synchronous cancers between the patients from BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “bilateral synchronous” which allows the options of yes and no.

Definitions:

Bilateral synchronous cancers are the cancers that occur in both breasts simultaneously or sequentially within three months of time frame.

2f: Menopausal status for invasive tumour by referral source[†]

Referral source	Pre	Post	Peri
BSA (N=645)	19.07%	67.29%	13.64%
Not BSA (N=1176)	31.63%	60.29%	8.08%
P value	< 0.001	0.003	< 0.001

[†] Referral source was not known for 4 invasive records. Menopausal status was not known for 13 BSA females and 17 not BSA females. There were 13 males in the not BSA group.

Comments:

The majority (63%) of the New Zealand patients were post-menopausal. The percentage of pre-menopausal women was significantly lower in the BSA group (19%) than in the not BSA group (32%). The BSA group had significantly higher percentages of post and peri-menopausal patients.

Audit data used:

Information is derived from the audit question “menopausal status” which allows the options of pre, peri, post and male.

Definitions:

Pre – an individual who has not yet experienced the menopause

Post – an individual who has experienced the menopause and the occurrence of greater than one year of spontaneous amenorrhoea

Peri – an individual who is either in the period just prior to the menopause or the subsequent one year of amenorrhoea following the menopause

Male – male patient

2g: Oestrogen receptor status of invasive tumour by referral source[†]

Referral source	Positive	Negative
BSA (N=642)	90.03%	9.97%
Not BSA (N=1180)	80.08%	19.92%
P value	<0.001	<0.001

[†] Referral source was not known for 4 invasive records. Oestrogen receptor status was not known for 16 BSA and 26 not BSA patients.

Comments:

Most (84%) of the New Zealand patients had oestrogen positive tumours. The percentage of patients with oestrogen positive tumours was significantly higher in the BSA group (90%) than in not BSA group (80%).

Audit data used:

Information is derived from the audit question “Oestrogen receptor status” which allows the options of positive, negative, ordered but not known and not done.

Definitions:

The presence or absence of oestrogen receptors on the tumour cells.

2h: Progesterone receptor status of invasive tumour by referral source[†]

Referral source	Positive	Negative
BSA (N=640)	78.59%	21.41%
Not BSA (N=1175)	65.02%	34.98%
P value	< 0.001	< 0.001

[†]Referral source was not known for 4 invasive records. Progesterone receptor status was not known for 18 BSA and 31 not BSA patients.

Comments:

The majority (70%) of the New Zealand patients had progesterone positive tumours. The percentage of patients with progesterone positive tumours was significantly higher in the BSA group (79%) than in not BSA group (65%).

Audit data used:

Information is derived from the audit question “progesterone receptor status” which allows the following options: positive, negative, ordered but not known and not done.

Definitions:

The presence or absence of progesterone receptors on the tumour cells.

2i: HER2 Receptor status of invasive tumour by referral source†

Referral source	Positive	Negative
BSA (N= 625)	10.40%	89.60%
Not BSA (N=1107)	18.16%	81.84%
P value	<0.001	<0.001

† Referral source was not known for 4 invasive records. HER2 status was not known for 33 BSA and 99 not BSA patients.

Comments:

Most (85%) of the New Zealand patients had HER2 negative invasive tumours. The percentage of patients with HER2 negative tumours was significantly higher in the BSA group (90%) than in not BSA group (82%).

Audit data used:

Information is derived from the audit question “HER2 receptor status” which allows the following options: positive, negative, ordered but not known and not done.

Definitions:

HER2 stands for **H**uman **E**pidermal growth factor **R**eceptor 2

Positive – biopsy revealed abnormally high levels of the HER2 gene or protein

Negative – biopsy revealed a normal level of the HER2 gene or protein

3. DCIS Tumour Characteristics

3a: Size of DCIS tumours by referral source[†]

Referral source	<10 mm	10-14 mm	15-19 mm	20-29 mm	30-39 mm	>40 mm
BSA (N=164)	34.15%	14.02%	11.59%	11.59%	10.37%	18.29%
Not BSA (N=79)	29.11%	11.39%	8.86%	22.78%	6.33%	21.52%
P value	0.433	0.570	0.520	0.023	0.304	0.551

[†] Referral source was not known for 3 DCIS records. DCIS tumour size was not known for 2 BSA and 9 not BSA patients.

Comments:

The percentage of patients with smaller tumours (<20mm) was higher for the BSA group (60%) than the not BSA group (40%) but this difference was not statistically significant.

Audit data used:

Information is derived from the audit question “DCIS tumour size in mm”.

Definitions:

Tumour size refers to the maximum diameter in millimetres of the furthest points of extension of the DCIS tumour cells in the principal tumour.

3b: Histological grade of DCIS tumour by referral source[†]

Referral source	Low	Intermediate	High
BSA (N=164)	9.76%	39.02%	51.22%
Not BSA (N=84)	8.33%	42.86%	48.81%
P value	0.715	0.560	0.719

[†] Referral source was not known for 3 DCIS records. DCIS Histological grade was not known for 2 BSA and 4 not BSA patients.

Comments:

The histological grade of the DCIS tumours was not significantly different between BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “DCIS histological grade of tumour” which allows the following options: low, medium and high.

Definitions:

The degree of differentiation of the breast cancer or the degree to which it resembles normal tissue as assessed by the pathologist.

Low – Well differentiated

Intermediate – Moderately differentiated

High – Poorly differentiated

3c: Necrosis of DCIS tumour by referral status[†]

Referral source	Absent	Present
BSA (N=159)	32.08%	67.92%
Not BSA (N=72)	31.94%	68.06%
P value	0.984	0.984

[†] Referral source was not known for 3 DCIS records. Necrosis of the DCIS tumours was not known for 7 BSA and 16 not BSA patients.

Comments:

The majority (68%) of New Zealand patients with DCIS tumours had necrosis. The percentage of DCIS tumours with necrosis was not significantly different between BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “necrosis of tumour” which allows the options of present and absent.

Definitions:

Two categories of necrosis are recognised with DCIS: focal necrosis with no central necrosis and central necrosis in ducts.

Present – Central necrosis is identified in ducts (this has previously been described as “comedo” type necrosis.).

Absent – Necrosis is not present or minimal. No central duct necrosis is present, but focal necrosis and isolated apoptotic cells may be present.

3d: Bilateral synchronous status of DCIS tumours by referral status[†]

Referral source	Bilateral synchronous	Not bilateral synchronous
BSA (N=166)	2.41%	97.59%
Not BSA (N=87)	6.90%	93.10%
P value	0.082	0.082

[†] Referral source was not known for 3 DCIS records. Bilateral synchronous status was not known for 1 not BSA patient.

Comments:

DCIS tumours in most (96%) of the New Zealand patients were not bilateral synchronous. The percentage of patients with bilateral synchronous DCIS tumours was lower in the BSA group (2%) than in not BSA group (7%) but this difference was not statistically significant.

Audit data used:

Information is derived from the audit question “bilateral synchronous” which allows the option of yes and no.

Definitions:

Bilateral synchronous cancers are cancers that occur in both breasts simultaneously or sequentially within three months.

3e: Menopausal status for the DCIS tumours by referral source[†]

Referral source	Pre	Post	Peri
BSA (N=159)	28.93%	61.64%	9.43%
Not BSA (N=87)	39.08%	57.47%	3.45%
P value	0.104	0.524	0.085

[†] Referral source was not known for 3 DCIS records. Menopausal status was not known for 7 BSA and 1 not BSA patients. There was 1 male from not BSA referral sources.

Comments:

The majority (60%) of the New Zealand DCIS patients were post-menopausal. The distribution of menopausal status was not significantly different between BSA and not BSA groups.

Audit data used:

Information is derived from the audit question “menopausal status” where the options are: pre, peri, post and male.

Definitions:

Pre – an individual who has not yet experienced the menopause

Post – an individual who has experienced the menopause and the occurrence of 12 months of spontaneous amenorrhoea

Peri – an individual who is either in the period just prior to the menopause or the subsequent 1 year of amenorrhoea following the menopause

Male – Male patient

4. Breast Surgery Treatment

4a: First breast surgery performed for invasive cancer by referral source[†]

Referral source	None	Open Biopsy	CLE	Mastectomy	Other
BSA (N= 658)	0.76%	2.89%	66.11%	29.18%	1.06%
Not BSA (N=1202)	5.74%	1.91%	35.11%	56.57%	0.67%
P value	<0.001	0.176	<0.001	<0.001	0.358

[†] Referral source was not known for 4 invasive records. Surgery status was not known for 4 not BSA participants.

Comments:

The majority of the BSA patients (69%) had breast conserving surgery (CLE or Open Biopsy) and the majority of not BSA patients had mastectomy (57%) as their first breast surgery.

4b: Further breast surgery after breast conserving surgery for invasive cancer by referral source[†]

Referral source (for invasive tumours with BCS surgery)	Mastectomy	Re-excision	Other surgery	Any further surgery	No further breast surgery
BSA (N=454)	10.13%	9.69%	3.08%	20.04%	79.96%
Not BSA (N=445)	16.18%	13.71%	4.04%	30.34%	69.66%
P value	0.007	0.061	0.551	<0.001	<0.001

[†] Referral source was not known for 4 invasive tumours. Surgery status was not known for 4 not BSA participants.

Please note that some of the patients had more than one surgery after BCS and therefore the percentage of any further surgery after BCS does not equal to the sum of the percentages of mastectomy, re-excision and other surgery after BCS.

Comments:

The majority (75%) of New Zealand patients had no further surgery after breast conserving surgery for invasive cancer. The percentage of patients undergoing mastectomy after BCS for invasive cancer was significantly less for BSA (10%) compared to not BSA groups (16%).

4c: Reconstruction after mastectomy for invasive cancer by referral source[†]

Referral source (for invasive tumours with mastectomy)	Reconstruction	No Reconstruction
BSA (N= 238)	14.29%	85.71%
Not BSA (N=752)	11.77%	88.23%
P value	0.304	0.304

[†] Referral source was not known for 3 invasive tumours which had mastectomy.

Comments:

The majority (88%) of New Zealand patients had no reconstruction after mastectomy for invasive cancer. The percentage of patients with reconstruction surgery after mastectomy for invasive tumours was not significantly different between BSA and not BSA groups.

4d: First breast surgery performed for DCIS by referral source†

Referral source	None	Open biopsy	CLE	Mastectomy	Other
BSA (N=166)	0.00%	8.43%	70.48%	20.48%	0.60%
Not BSA (N=88)	4.55%	12.50%	50.00%	32.95%	0.00%
P value	NC	0.301	<0.001	0.029	NC

† Referral source was not known for 3 DCIS records.

Comments:

The majority (67%) of New Zealand patients had breast conserving surgery (open biopsy or CLE) as their first surgery for DCIS. The percentage of patients with breast conserving surgery (open biopsy or CLE) for DCIS tumours was significantly higher in the BSA group (79%) than in the not BSA group (63%). The percentage of patients with mastectomy as their first surgery for DCIS tumours was significantly lower in the BSA group (20%) than in the not BSA group (33%).

4e: Further surgery after breast conserving surgery for DCIS by referral source†

Referral source (for DCIS tumours with BCS surgery)	Mastectomy	Re-excision	Other	Any further surgery	No further breast surgery
BSA (N=131)	14.50%	22.90%	7.25%	38.16%	61.84%
Not BSA (N=55)	23.64%	21.82%	9.83%	45.45%	54.55%
P value	0.132	0.872	0.798	0.355	0.355

† Referral source was not known for 3 DCIS records. Please note that some of the patients had more than one surgery after BCS and therefore the percentage of any further surgery after BCS does not equal to the sum of the percentages of mastectomy, re-excision and other surgery after BCS.

Comments:

The majority (60%) of New Zealand patients had no further surgery after breast conserving surgery for DCIS. The percentage of patients with mastectomy after breast conserving surgery for DCIS tumours was lower in the BSA group (15%) than in the not BSA group (24%) but this difference was not significant.

4f: Reconstruction performed after mastectomy for DCIS by referral source†

Referral source (for DCIS tumours with mastectomy)	Reconstruction	No reconstruction
BSA (N=53)	32.08%	67.92%
Not BSA (N=42)	42.86%	57.14%
P value	0.279	0.279

† Referral source was not known for 3 DCIS records.

Comments

The majority (63%) of the New Zealand patients had no reconstruction after mastectomy for DCIS. The percentage of patients with reconstruction surgery after mastectomy for DCIS was higher in the not BSA group (43%) compared with the BSA group (32%) but this difference was not significant.

Audit data used:

Information is derived from the audit question “surgical procedures” which allows the following options: no surgery, ABBI, open biopsy, CLE, re-excision, total mastectomy, reconstruction and other.

Definitions:

Open biopsy (including localisation) - Surgical procedure in which a sample of breast tissue for histological examination is obtained in a conventional surgical procedure, using an open excision.

CLE - The complete excision of an entire tumour mass.

ABBI – The process whereby an Advanced Breast Biopsy Instrumentation System (or similar) technique is used to excise non-palpable breast lesions.

Total mastectomy - The surgical removal of the breast

Re-excision – A secondary surgical procedure conducted to obtain a rim of normal breast tissue around the periphery of the previously removed primary tumour.

Reconstruction – The use of a prosthesis or tissue from other parts of the body to re-build a breast.

Other – Other surgery

5. Axillary surgery treatment

Relevant clinical practice guidelines

- Women with unifocal ≤ 3 cm invasive tumours and clinically negative nodes should be offered sentinel node biopsy^{1,3}.
- For women with multifocal >3 cm invasive tumours with clinically involved nodes, axillary dissection is normally recommended^{1,3}.
- Axillary dissection should not be performed in the management of DCIS unless invasion is suspected^{1,2}.

NBCA KPIs

- KPI 3 - Percent of cases undergoing axillary surgery for invasive cancer ($>90\%$).
- KPI 4: Percent of cases not undergoing axillary surgery for DCIS which underwent breast conserving surgery ($>90\%$).

5a. Axillary procedures for invasive cancer by referral source[†]

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 or 3 only	SNB & Level 1	SNB & Level 2 or 3	Level 1 & 2 or 3	Level 2 & 3	>2 axillary surgeries
BSA (N=652)	2.45%	65.95%	1.23%	15.18%	0.61%	13.80%	0.15%	0.46%	0.15%
Not BSA (N=1196)	8.28%	35.70%	0.84%	38.04%	0.33%	15.72%	0.42%	0.42%	0.25%
P value	<0.001	<0.001	0.414	<0.001	0.383	0.271	0.474	0.895	0.667

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 &/or 3	SNB & Level 1 only	SNB & Level 2 or 3
BSA (N=652)	2.45%	65.95%	1.23%	15.94%	0.61%	13.80%
Not BSA (N=1196)	8.28%	35.70%	0.84%	39.13%	0.33%	15.72%
P value	<0.001	<0.001	0.414	<0.001	0.383	0.271

[†] Referral source was not known for 4 invasive tumours. Axillary procedures was not known for 6 BSA and 10 not BSA patients.

5b. Axillary procedures for ≤ 3 cm invasive cancer by referral source[†]

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 or 3 only	SNB & Level 1	SNB & Level 2 or 3	Level 1 & 2 or 3	Level 2 & 3	>2 axillary surgeries
BSA (N=601)	2.00%	70.05%	1.33%	11.81%	0.67%	13.81%	0.17%	0.17%	0.00%
Not BSA (N=865)	3.35%	45.66%	1.04%	31.56%	0.23%	17.46%	0.35%	0.12%	0.23%
P value	0.121	< 0.001	0.609	< 0.001	0.200	0.061	0.515	0.796	Not Calculated

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 &/or 3	SNB & Level 1 only	SNB & Level 2 or 3
BSA (N=601)	2.00%	70.05%	1.33%	12.15%	0.67%	13.81%
Not BSA (N=865)	3.35%	45.66%	1.04%	32.26%	0.23%	17.46%
P value	0.121	< 0.001	0.609	<0.001	0.200	0.061

[†] For ≤ 3 cm tumours, referral source was not known for 3 patients, axillary procedure was not known for 2 BSA and 8 not BSA patients, tumour size was missing for 9 BSA and 72 not BSA patients. Most of the patients with missing tumour size information were recorded as having no axillary surgery.

5c. Axillary procedures for >3cm invasive cancer by referral source by referral source[†]

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 or 3 only	SNB & Level 1	SNB & Level 2 or 3	Level 1 & 2 or 3	Level 2 & 3	>2 axillary surgeries
BSA (N=45)	2.22%	15.56%	0.00%	60.00%	0.00%	15.56%	0.00%	4.44%	2.22%
Not BSA (N=260)	5.00%	11.15%	0.38%	66.15%	0.77%	14.23%	0.38%	1.54%	0.38%
<i>P value</i>	0.411	0.398	Not calculated	0.423	Not calculated	0.815	Not Calculated	0.195	0.159

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 &/or 3	SNB & Level 1 only	SNB & Level 2 or 3
BSA (N=45)	2.22%	15.56%	0.00%	66.66%	0.00%	15.56%
Not BSA (N=260)	5.00%	11.15%	0.38%	68.45%	0.77%	14.23%
<i>P value</i>	0.411	0.398	Not calculated	0.811	Not calculated	0.815

[†] For >3 cm tumours, referral source not known for one patient, axillary procedure was not known for 1 BSA and 1 not BSA patients, tumour size was missing for 9 BSA and 72 not BSA patients. Most of the patients with missing tumour size information were recorded as having no axillary surgery.

Comments: For ≤ 3 cm tumours, a higher percentage (70%) of BSA patients had SNB as their only axillary surgery compared with not BSA patients (46%). This difference was not significant for >3cm tumours. As expected from the guidelines, a higher percentage of patients had Level 2 or Level 3 axillary surgery (with or without SNB) for >3cm tumours (83%) than for ≤ 3 cm tumours (40%).

5d. Axillary procedures for DCIS treated with breast conserving surgery only by referral source†

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 or 3 only	SNB & Level 1	SNB & Level 2 or 3	Level 1 & 2 or 3	Level 2 & 3	>2 axillary surgeries
BSA (N=130)	83.08%	16.15%	0.00%	0.00%	0.77%	0.00%	0.00%	0.00%	0.00%
Not BSA (N=53)	64.15%	32.08%	3.77%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
P value	0.005	0.016				Not calculated			

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 &/or 3	SNB & Level 1 only	SNB & Level 2 or 3
BSA (N=130)	83.08%	16.15%	0.00%	0.00%	0.77%	0.00%
Not BSA (N=53)	64.15%	32.08%	3.77%	0.00%	0.00%	0.00%
P value	0.005	0.016			Not calculated	

†There were 1 BSA and 2 not BSA patients with not known axillary procedure for DCIS with BCS.

Comments:

The majority (78%) of New Zealand patients with DCIS did not have axillary surgery after BCS, as expected from the guidelines. Significantly less BSA patients (17%) had axillary surgery than not BSA patients (36%).

5e. Axillary procedures for DCIS treated with mastectomy by referral source[†]

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 or 3 only	SNB & Level 1	SNB & Level 2 or 3	Level 1 & 2 or 3	Level 2 & 3	>2 axillary surgeries
BSA (N=52)	30.77%	67.31%	0.00%	0.00%	1.92%	0.00%	0.00%	0.00%	0.00%
Not BSA (N=40)	22.50%	70.00%	5.00%	2.5%	0.00%	0.00%	0.00%	0.00%	0.00%
P value	0.377	0.783				Not calculated			

Referral source	No axillary surgery	SNB only	Level 1 only	Level 2 &/or 3	SNB & Level 1 only	SNB & Level 2 or 3
BSA (N=52)	30.77%	67.31%	0.00%	0.00%	1.92%	0.00%
Not BSA (N=40)	22.50%	70.00%	5.00%	2.5%	0.00%	0.00%
P value	0.377	0.783			Not calculated	

[†] There were 1 BSA and 2 not BSA patients with not known axillary procedure for DCIS with mastectomy

Comments:

The majority (73%) of New Zealand patients with DCIS had axillary surgery after mastectomy. Number of axillary surgery performed was not significantly different between the BSA group and not BSA group.

Audit data used:

Information on axillary procedure is from the audit question “axillary surgery” which allows the following options: sentinel node biopsy; level 1 sampling, level 2, level 3 and no axillary surgery.

Definitions:

Sentinel node biopsy - The identification and excision of the sentinel lymph node (the first node(s) draining the primary tumour in the regional lymphatic basin) from patients with invasive breast cancer.

Axillary surgery - The surgical excision of the axillary contents (fat and lymph nodes) *en bloc* with mastectomy or as an independent procedure.

Level 1 - The excision of a single, low axillary node or the excision of the axillary contents up to the inferior border of the pectoralis minor muscle, includes sampling.

Level 2 – Excision of the axillary contents up to the superior border of the pectoralis minor muscle.

Level 3 - Excision of the axillary contents up to the apex of the axilla.

6. Margins of excision for breast surgery

6a. Margins of excision for invasive cancer by referral source[†]

Referral source	Margin size =0 mm	Margin size > 0 mm and ≤ 1 mm	Margin size ≥ 2 mm
BSA (N=608)	2.80%	6.25%	90.95%
Not BSA (N=1073)	5.22%	11.09%	83.69%
P value	0.019	0.001	<0.001

[†] Referral source was not known for 4 invasive cases. Margin size was not known for 50 BSA and 133 not BSA patients.

Comments:

Most (86%) of the New Zealand patients had ≥2 mm margins after surgery for invasive cancer. The percentage of patients with ≥2 mm margins after the surgery for invasive tumour was significantly higher in the BSA group (91%) than in not BSA group (84%). The percentage of patients with involved margin after the surgery for invasive tumour was significantly lower in the BSA group (3%) than in not BSA group (5%).

6b. Margins of excision for DCIS cancer by referral source[†]

Referral source	Margin size =0 mm	Margin size > 0 mm and ≤ 1 mm	Margin size ≥ 2 mm
BSA (n=158)	5.06%	15.82%	79.11%
Not BSA (N=73)	9.59%	15.07%	75.34%
P value	0.194	0.883	0.521

[†] Margin size was not known for 8 BSA and 15 not BSA patients for DCIS cases.

Comments:

Most (78%) New Zealand patients had ≥2 mm margins after surgery for DCIS. Margin size was not significantly different between BSA and not BSA groups.

Audit data used:

Information on clear margin is derived from the audit question “distance (in mm) to closest circumferential margin” and “distance (in mm) to closest vertical margin” which allows the user entry of margin size in mm.

Definitions:

Margin size = lowest value recorded for vertical margin or circumferential margin.

7. Radiotherapy treatment

NBCA KPIs

- KPI 1: Percent of invasive tumours treated with breast conserving surgery (BCS) that were referred for or prescribed radiotherapy ($\geq 85\%$).
- KPI 4: High risk mastectomy cases that were referred for or prescribed radiotherapy. High risk defined as: invasive tumour size ≥ 50 mm OR invasive tumours with ≥ 4 positive lymph nodes ($\geq 85\%$).

7a: Radiotherapy for invasive cancer treated with breast conserving surgery by referral source[†]

Referral source	Radiotherapy prescribed	Radiotherapy not prescribed
BSA (N=390)	96.92%	3.08%
Not BSA(N=343)	96.21%	3.79%
P value	0.596	0.596

[†] Referral source was not known for 1 BCS invasive cases. Radiotherapy was not known for 14 BSA and 26 not BSA patients. Please note that the patients who had mastectomy or other breast surgery after breast conserving surgery were not included in this group.

Comments:

The percentage of patients prescribed with radiotherapy treatment after breast conserving surgery for invasive cancer was not significantly different between the BSA (97%) and not BSA groups (96%).

7b: Radiotherapy for invasive cancer which had mastectomy by referral source[†]

Referral source	Radiotherapy prescribed	Radiotherapy not prescribed
BSA (N=228)	33.33%	66.67%
Not BSA (N=717)	48.68%	51.32%
P value	< 0.001	< 0.001

[†] Referral source was not known for 3 invasive mastectomy cases and radiotherapy status was not known for 10 BSA and 35 not BSA patients.

Comments:

The percentage of patients with prescribed radiotherapy treatment after mastectomy for invasive cancer was significantly lower in the BSA group (33%) than in the not BSA group (49%).

7c: Radiotherapy for high risk invasive cancer treated with mastectomy by referral source[†]

Referral source	Radiotherapy prescribed	Radiotherapy not prescribed
BSA (N=30)	90.00%	10.00%
Not BSA (N=200)	87.50%	12.50%
P value	0.696	0.696

[†] There were 9 patients from not BSA referral sources with not known radiotherapy data.

Comments:

The percentage of patients with prescribed radiotherapy treatment after mastectomy for high risk invasive cancer was not significantly different between BSA group and not BSA group. The percentage of patients receiving radiotherapy in high risk mastectomy group is much higher than in the whole mastectomy group.

7d: Radiotherapy for DCIS treated with breast conserving surgery by referral source[†]

Referral source	Radiotherapy prescribed	Radiotherapy not prescribed
BSA (N=109)	81.65%	18.35%
Not BSA (N=39)	69.23%	30.77%
P value	0.106	0.106

[†] Radiotherapy status was not known for 2 BSA and 3 not BSA patients. Please note that the DCIS patients who had mastectomy after breast conserving surgery were excluded in this group.

Comments:

The majority (78%) of New Zealand patients had radiotherapy after breast conserving surgery. There was no significant difference in radiotherapy treatment after breast conserving surgery for DCIS between the BSA and not BSA groups.

7e: Radiotherapy for DCIS cancer which had mastectomy by referral source[†]

Referral source	Radiotherapy prescribed	Radiotherapy not prescribed
BSA (N=49)	6.12%	93.88%
Not BSA (N=37)	5.41%	94.59%
P value	0.888	0.888

[†] Radiotherapy status was not known for 4 BSA and 5 not BSA patients.

Comments:

Only a small percentage (6%) of New Zealand patients had radiotherapy treatment after mastectomy for DCIS cancer. There was no significant difference in radiotherapy treatment after mastectomy for DCIS between the BSA and not BSA groups.

Audit data used:

Information on patients undergoing radiotherapy is derived from the audit question “did you prescribe or refer for any of the following adjuvant therapies?” where one of the options is radiotherapy. The options were: yes, no, not yet, referred but not used, not known.

Definitions:

Radiotherapy is the use of radiation, usually X-rays or gamma rays, to kill tumour cells.

8. Hormonal treatment

NBCA KPIs⁵

- KPI 2: Percent referred or prescribed hormonal treatment for oestrogen positive invasive tumours ($\geq 85\%$).

8a. Hormonal treatment for oestrogen positive invasive tumours by referral source[†]

Referral source	Hormonal treatment	No hormonal treatment
BSA (N=540)	85.37%	14.63%
Not BSA (N=850)	90.71%	9.29%
P value	0.002	0.002

Referral source	Menopausal status	Any hormone treatment		Tamoxifen & Ovarian ablation		Ovarian ablation & Aromatase inhibitors		All three hormone treatments		No hormone treatment
		Any hormone treatment	Tamoxifen only	Tamoxifen & Ovarian ablation	Tamoxifen & Aromatase inhibitors	Ovarian ablation only	Ovarian ablation & Aromatase inhibitors	Aromatase inhibitors only	All three hormone treatments	No hormone treatment
BSA (N=540)	Any	85.37%	45.00%	0.00%	9.81%	0.37%	0.00%	30.19%	0.00%	14.63%
Not BSA (N=850)	Any	90.71%	50.59%	0.47%	8.00%	0.24%	0.00%	31.06%	0.35%	9.29%
P value		0.002	0.042	NC	0.242	0.647	NC	0.731	NC	0.002
BSA (N=112)	pre	87.50%	75.89%	0.00%	6.25%	1.79%	0.00%	3.57%	0.00%	12.50%
Not BSA (N=253)	pre	94.86%	85.38%	1.19%	4.35%	0.79%	0.00%	2.37%	0.79%	5.14%
P value		0.013	0.028	NC	0.439	0.400	NC	0.517	NC	0.013
BSA (N=72)	peri	86.11%	63.89%	0.00%	11.11%	0.00%	0.00%	11.11%	0.00%	13.89%
Not BSA (N=67)	peri	91.04%	62.69%	0.00%	11.94%	0.00%	0.00%	16.42%	0.00%	8.96%
P value		0.362	0.883	NC	0.878	NC	NC	0.363	NC	0.362
BSA (N=350)	post	85.71%	32.00%	0.00%	10.86%	0.00%	0.00%	42.86%	0.00%	14.29%
Not BSA (N=515)	post	88.93%	31.07%	0.19%	9.51%	0.00%	0.00%	47.96%	0.19%	11.07%
P value		0.158	0.772	NC	0.519	NC	NC	0.139	NC	0.158
BSA (N=0)	Male	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
Not BSA (N=10)	Male	100.00%	100.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
P value		NC	NC	NC	NC	NC	NC	NC	NC	NC

BSA (N=6)	unknown	16.67%	0.00%	0.00%	0.00%	0.00%	0.00%	16.67%	0.00%	83.33%
Not BSA (N=5)	unknown	40.00%	40.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	60.00%
P value		0.387	NC	NC	NC	NC	NC	NC	NC	0.387

† Referral source was not known for 1 record. Oestrogen receptor status was not known for 16 BSA records and 26 not BSA records. Hormonal treatment was not known for 38 BSA and 95 not BSA patients for oestrogen positive tumours.

Comments:

Most (89%) of the New Zealand patients with oestrogen positive tumours had hormonal treatment. The percentage of patients with prescribed hormonal treatment for oestrogen positive invasive tumours was significantly lower in the BSA group (85%) than in not BSA group (91%).

8b. Hormonal treatment for oestrogen negative invasive cancer by referral source[†]

Referral source	Hormonal treatment	No hormonal treatment
BSA (N=59)	11.86%	88.14%
Not BSA (N=219)	6.39%	93.61%
P value	0.158	0.158

Referral source	Menopausal status	Any hormone treatment		Tamoxifen & Ovarian ablation	Tamoxifen & Aromatase inhibitors	Ovarian ablation only	Ovarian ablation & Aromatase inhibitors	Aromatase inhibitors only	All three hormone treatments	No hormone treatment
		Any hormone treatment	Tamoxifen only	Tamoxifen & Ovarian ablation	Tamoxifen & Aromatase inhibitors	Ovarian ablation only	Ovarian ablation & Aromatase inhibitors	Aromatase inhibitors only	All three hormone treatments	No hormone treatment
BSA (N=59)	Any	11.86%	5.08%	0.00%	1.69%	1.69%	0.00%	1.69%	1.69%	88.14%
Not BSA (N=219)	Any	6.39%	5.02%	0.00%	0.46%	0.00%	0.00%	0.91%	0.00%	93.61%
P value		0.158	0.985	NC	0.318	NC	NC	0.606	NC	0.158
BSA (N=4)	pre	25.00%	25.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	75.00%
Not BSA (N=71)	pre	5.63%	5.63%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	94.37%
P value		0.131	0.131	NC	NC	NC	NC	NC	NC	0.131
BSA (N=8)	peri	25.00%	0.00%	0.00%	0.00%	0.00%	0.00%	12.50%	12.50%	75.00%
Not BSA (N=21)	peri	9.52%	9.52%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	90.48%
P value		0.280	NC	NC	NC	NC	NC	NC	NC	0.280
BSA (N=45)	post	8.89%	4.44%	0.00%	2.22%	2.22%	0.00%	0.00%	0.00%	91.11%
Not BSA (N=123)	post	5.69%	3.25%	0.00%	0.81%	0.00%	0.00%	1.63%	0.00%	94.31%
P value		0.458	0.712	NC	0.456	NC	NC	NC	NC	0.458
BSA (N=0)	Male	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
Not BSA (N=2)	Male	50.00%	50.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	50.00%
P value		NC	NC	NC	NC	NC	NC	NC	NC	NC
BSA (N=2)	UK	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	100.00%
Not BSA (N=2)	UK	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	100.00%
P value		NC	NC	NC	NC	NC	NC	NC	NC	1.00

[†] Referral source was not known for 4 invasive records. Oestrogen receptor status was not known for 16 BSA and 26 not BSA patients. Hormonal treatment was not known for 5 BSA and 16 not BSA patients for oestrogen negative invasive tumours.

Comments:

A small percentage (8%) of New Zealand patients with oestrogen negative tumours had hormonal treatment. These findings were not significantly different between BSA and not BSA groups.

Audit data used:

Information for oestrogen receptor positive status is derived from the audit question “receptor status” where information is recorded for oestrogen and progesterone status with options of positive, negative, ordered but not known and not done.

Information for number of patients prescribed and/or referred for hormonal therapy is derived from the question “did you prescribe or refer for any of the following adjuvant therapies?” The following options apply: yes, no, not yet and referred but not used.

Definitions:

Oestrogen receptors are prognostic indicators. They are an intracellular receptor protein that binds oestrogens and anti-oestrogens, mediating their effects by binding to DNA and altering the expression of specific genes.

Hormonal treatment – SERMs, aromatase inhibitors, ovarian ablation

SERMSs- The use of Selective Oestrogen Receptor Modulators to inhibit the growth of hormone responsive cancer cells after primary treatment, either by surgery or radiotherapy or a combination of these, to eradicate micro metastatic cancer.

Ovarian ablation – The use of surgery, radiation or drug treatment to cease hormone production by the ovaries, after primary treatment either by surgery or radiotherapy or a combination of these (usually within six weeks), to eradicate micro metastatic cancer.

Aromatase inhibitors – These are a class of drugs which lower the level of oestrogen in the tumour. They are primarily used in post-menopausal patients.

9. Chemotherapy treatment

9a. Chemotherapy treatment for invasive cancer for ≤ 70 years old patients by referral source[†]

Referral source	Chemotherapy prescribed	Chemotherapy not prescribed
BSA (N=618)	38.83%	61.17%
Not BSA (N=772)	71.37%	28.63%
<i>P value</i>	< 0.001	< 0.001

Referral source	Chemotherapy prescribed			Chemotherapy not prescribed		
	Oestrogen/Progesterone			Oestrogen/Progesterone		
	Positive	Negative	Unknown	Positive	Negative	Unknown
BSA (N=618)	31.39%	6.80%	0.65%	57.28%	2.75%	1.13%
Not BSA (N=772)	53.11%	17.88%	0.39%	26.04%	1.68%	0.91%
<i>P value</i>	<0.001	<0.001	0.498	<0.001	0.174	0.675

[†] Referral source was not known for 4 records. Chemotherapy status was not known for 24 BSA and 61 not BSA patients

Comments:

The percentage of ≤ 70 years old patients with prescribed chemotherapy treatment was significantly lower in BSA group (39%) than in not BSA group (71%).

9b. Chemotherapy treatment for invasive cancer for > 70 years old patients by referral source[†]

Referral source	Chemotherapy prescribed	Chemotherapy not prescribed
BSA (N=13)	7.69%	92.31%
Not BSA (N=361)	17.17%	82.83%
<i>P value</i>	<0.001	<0.001

Referral source	Chemotherapy prescribed			Chemotherapy not prescribed		
	Oestrogen/Progesterone			Oestrogen/Progesterone		
	Positive	Negative	Unknown	Positive	Negative	Unknown
BSA (N=13)	7.69%	0.00%	0.00%	92.31%	0.00%	0.00%
Not BSA (N=361)	9.14%	8.03%	0.00%	72.02%	9.14%	1.66%
<i>P value</i>	0.858	NC	NC	0.107	NC	NC

[†] Chemotherapy status was not known for 3 BSA and 12 not BSA patients.

Comments:

A small percentage (17%) of >70 year old New Zealand patients had chemotherapy treatment. The percentage of >70 years old patients prescribed chemotherapy treatment was significantly lower in the BSA group (8%) than in the not BSA group (17%).

Audit data used:

Information on chemotherapy was derived from the audit question “did you prescribe or refer for any of the following adjuvant therapies?” where one choice is chemotherapy. The following options apply: yes, no, not yet, referred but not used.

Definitions:

Chemotherapy is the use of cytotoxic drugs that aim to kill, prevent or slow the growth rate of cancer cells.

10. Herceptin (Trastuzumab) treatment

Relevant Clinical Practice Guidelines

Patients with early breast cancer and HER-2 positive tumours, either node positive or node negative with tumours larger than 1cm, should be offered trastuzumab with chemotherapy following surgery⁴.

10a. Herceptin (Trastuzumab) treatment for >1cm HER2 positive OR node positive HER2 positive invasive cancer by referral source[†]

Referral source	Herceptin prescribed			Herceptin not prescribed
	Chemo yes	Chemo no	Chemo unknown	
BSA (N=46)	86.96%	0.00%	0.00%	13.04%
Not BSA (N=166)	75.90%	3.01%	1.20%	19.88%
P value	0.108	Not calculated		0.290

[†] Referral source was not known for 4 invasive records. Herceptin treatment was not known for 4BSA patients and 10 not BSA patients with >1cm HER2 positive or node positive HER2 positive tumours.

Comments:

The majority (78%) of the New Zealand patients with HER2 positive >1 cm or node positive tumours received Herceptin treatment. There was no significant difference between the BSA and not BSA patients regarding Herceptin treatment for these patients.

Audit data used:

Information on chemotherapy was derived from the audit question “did you prescribe or refer for any of the following adjuvant therapies?” where one choice is Herceptin or other immunotherapy. The following options apply: yes, no, not yet, referred but not used.

Definitions:

Herceptin is a drug aimed at women who show HER2 gene amplification and/or protein over expression.

11. Summary

While the majority of breast cancer cases from New Zealand were referred as symptomatic from a GP (52%), BSA was the second most common referral source for New Zealand breast cancer patients (39%).

The majority (88%) of New Zealand cases were invasive breast cancer. The percentage of DCIS cancer was higher in the BSA group (20%) compared to not BSA group (7%).

The majority of New Zealand patients were public (72%). The percentages of public and private patients were not significantly different between the BSA group and not BSA group.

The percentage of 51 -70 year old patients was higher in the BSA group (76%) than in the not BSA group (36%). The percentage of younger (< 40 years) and older patients (> 70 years) were lower in the BSA group (0.1%, 3%) compared with the not BSA group (9%, 31%). The percentage of patients in the 41-50 years group was not significantly different between the BSA and not BSA groups.

Only 1% of the New Zealand patients were males and none of the male patients were referred from BSA for the treatments.

There were some differences between BSA and not BSA patients for the invasive and DCIS tumour characteristics and accordingly there were differences in some of the breast cancer treatments between BSA and not BSA patients (Table 11a and Table 11b).

11a. Significant differences in invasive tumour characteristics and breast cancer treatments between BSA and not BSA patients

Invasive tumour characteristics that were significantly different between BSA and not BSA patients

Higher percentage of BSA patients (3%) had Invasive Tubular tumours compared to not BSA patients (1%).

Higher percentage of BSA patients (57%) had smaller (<15 mm) tumours compared to not BSA patients (24%).

Lower percentage of BSA patients (24%) had larger tumours (>20 mm) compared to not BSA patients (60%).

Higher percentage of BSA patients (35%) had invasive Grade 1 tumours compared to not BSA patients (18%).

Lower percentage of BSA patients (19%) had Grade 3 tumours compared to not BSA patients (40%).

Lower percentage of BSA patients (14%) had lymphatic vascular invasion compared to not BSA patients (34%).

Lower percentage of BSA patients (19%) were pre-menopausal compared not BSA patients (32%)

Higher percentage of BSA patients (90%) had oestrogen positive tumours compared to not BSA patients (80%).

Higher percentage of BSA patients (79%) had progesterone positive tumours compared to not BSA patients (65%).

Lower percentage of BSA patients (10%) had HER2 positive tumours compared to not BSA patients (18%).

Treatments for invasive tumours that were significantly different between BSA and not BSA patients

Breast surgery

The majority of BSA patients (69%) had BCS as their first breast surgery. The majority of not BSA patients had mastectomy as their first breast surgery (57%).

Further surgery after breast conserving surgery

A smaller percentage of BSA patients (20%) had further surgery after breast conserving surgery compared with not BSA patients (30%).

Axillary surgery

For ≤ 3 cm tumours, a higher percentage (70%) of BSA patients had SNB as their only axillary surgery compared with not BSA patients (46%).

Margin after surgery

A higher percentage of BSA patients (91%) had ≥ 2 mm margins compared to not BSA patients (84%).

Radiotherapy treatment after mastectomy

A lower percentage of BSA patients (33%) were prescribed radiotherapy after mastectomy compared to not BSA patients (49%).

Hormone treatment

A lower percentage of BSA patients (85%) were prescribed hormonal treatment compared to not BSA patients (91%).

Chemotherapy treatment

The percentage of ≤ 70 years old patients with prescribed chemotherapy treatment was significantly lower in BSA group (39%) than in not BSA group (71%).

11b. DCIS tumour characteristics and breast cancer treatments between BSA and not BSA patients

DCIS tumour characteristics that were significantly different between BSA and not BSA patients

None of the DCIS characters were significantly different between BSA and not BSA groups.

Treatments for DCIS tumours that were significantly different between BSA and not BSA patients

Breast Surgery

A higher percentage of BSA patients (79%) had breast conserving surgery (CLE or open biopsy) as their first breast surgery compared to not BSA patients (63%).

The percentage of patients with mastectomy as their first surgery for DCIS tumours was significantly lower in the BSA group (20%) than in the not BSA group (33%).

Axillary Surgery

A smaller percentage of BSA patients (17%) had axillary surgery after BCS compared to not BSA patients (36%).

References

1. New Zealand Guidelines Group (NZGG). Management of Early Breast Cancer. Evidence-based Best Practice Guideline. First edition, New Zealand, 2009.
2. National Breast Cancer Centre. The clinical management of ductal carcinoma in situ, lobular carcinoma in situ and atypical hyperplasia of the breast: First edition, Camperdown, NSW, 2003
3. National Breast and Ovarian Cancer Centre. Recommendations for the use of sentinel node biopsy in early (operable) breast cancer. Canberra, June 2008
4. National Breast Cancer Centre. Recommendations for the use of trastuzumab (Herceptin®) for the treatment of HER2-positive breast cancer. Canberra: Commonwealth of Australia, 2007.